

# HNV farming in France

**Xavier POUX, EFNCP-AScA**





# An overview of French agrarian geography

- 🦒 **France at the crossroad of the main biogeographic zones**
- 🦒 **A great variety of geographical zones**
  - ✓ mountains, hills, plains
  - ✓ X humid and dry
- 🦒 **A great variety of « terroirs » inherited from the XIXth century**




# An overview of French goeography for EU citizens (in 1 mn)


Petites régions  
agricoles




06/10/1999



# When a large part of French farming systems used to be HNV



# Map of rural landscapes in 1950





# Main trends affecting French agriculture



## Modernisation of agriculture

- ✓ More fertilisers
- ✓ More plant protection products
- ✓ More tractors
- ✓ Less farmers !



## Saltus decreases (-30 % permanent grassland between 1970 / 2006)



- ✓ Fertilised permanent grassland
- ✓ Forage maize
- ✓ Cultivated grassland
- ✓ More crop




## Specialisation of farming systems

- ✓ Crop system
- ✓ Grass-based systems
- ✓ Imported protein crops (soyabean)






# Price index since 1960





# Development of French UAA since 1989




 **1950 : permanent grassland fertilised 8**


**1982 : permanent grassland fertilised 40 %**

**1998 : permanent grassland fertilised 66 %**


## Development of UAA and permanent grassland index (1979 = 100)



● **Chargement : total UGB  
herbivore/ha SFP**




● **Evolution de 1988 à 2000**



source: © Agreste – Recensements agricoles 1988 et 2000, traitement et cartographie: Institut de l'Elevage



## Likely density of type 1 HNV farming systems



### Légende


#### Chargement en bétail:

- < 0,7 UGB/ha en zone méditerranéenne ;
- < 1 UGB/ha en zone continentale et alpine ;
- compris entre 1 et 1,15 UGB/ha en zone atlantique.
- < 1 UGB/ha en zone méditerranéenne ;
- compris entre 1 et 1,15 UGB/ha en zone continentale.

#### Pacages collectifs

- Plus de 10 fermes par commune utilisant des pacages collectifs

## Likely density of type 2 HNV farming systems



### Légende

#### Chargement en bétail :

- < 1 UGB/ha en zone méditerranéenne, alpine et continentale ;
- compris entre 1 et 1,15 UGB/ha en zone atlantique.
- compris entre 1 et 1,15 UGB/ha en zone continentale.

#### Vergers traditionnels:

- densité par commune (> 4 pourcents de la SAU)

# Type 2 HNV : the most drastic decrease in the last decades

- ❖ Mixed farming is the most affected
- ❖ Loss of permanent grassland is the most visible in the mixed regions
- ❖ Low intensity dairy production hardly resists in such conditions
- ❖ What does happen when you go from 3000 kg milk/cow to 7000 kg?
- ❖ Permanent grassland are then intensified





● **Chargement : total UGB**


Livestock : =  
forage : -

Livestock : -  
forage : - -



0 75 150  
Kilomètres


● **Evolution de 1988 à 2000**



source: © Agreste – Recensements agricoles 1988 et 2000, traitement et cartographie: Institut de l'Elevage



# Pays d'Auge








# What does happen to type 1 HNV?

- 🐄 **Still large share of permanent grassland in France**
- 🐄 **But in detail:**
  - ✓ Intensification
  - ✓ Abandonment
- 🐄 **An insidious shift**
- 🐄 **Towards type 2?**






Mostly intensification  
(dairy and beef)

source: © Agres

Black & white:  
Abandonment vs. intensification  
(dairy and sheep)

● Evolution de 1988 à 2000



stitut de l'Elevage





Vercors, where permanent grassland is 98% of UAA in 2000



# Type 3

- ❖ In practice: mainly N2000 sites for birds
- ❖ The example of the little bustard (*tetrax tetrax*): rare and endangered bird with hardly no SNV
- ❖ Is that still HNV?
- ❖ Can it support other components of biodiversity (including saltus)?









# Conclusions

- ❖ **HNV farming is still rapidly decreasing in France**
- ❖ **Can we recover large share of type 1 HNV?**
  - ✓ Maintain the common pasture lands
- ❖ **Maps and monitoring should show the main threats (e.g. according to regions and productions)**
- ❖ **Needs to understand which farming systems should be supported**



# At landuse and landscape level

A bit of everything in each FS?  
( $\pm$  type 2)



HNV and LNV farming systems?



# The future of HNV farming in France

- ❗ **Very difficult as « extensification » is rejected by most parties (ex of Saône et Loire)**
  - ✓ Exception: Beaufort and, in a way Jura
- ❗ **What is accepted is a share of agro-ecological infrastructures (=SNV) in a more intensive context (type 2)**
- ❗ **Gouvernement is quite reluctant to HNV**
  - ✓ OK for « grass » and « permanent grassland »
  - ✓ Not for extensive grassland
- ❗ **Preferred approach: biodiversity in PDO**

